



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Sustained efficacy of natalizumab in the treatment of relapsing-remitting multiple sclerosis independent of disease activity and disability at baseline: real-life data from a Swiss cohort

Kallweit, U ; Jelcic, I ; Braun, N ; Fischer, H ; Zörner, B ; Schreiner, B ; Sokolov, A A ; Martin, R ; Weller, M ; Linnebank, M

Abstract: OBJECTIVES: Therapy for relapsing-remitting multiple sclerosis with natalizumab (Tysabri; Biogen Idec) has been shown to be effective in the reduction of the clinical relapse rate and disability progression. However, real-life longitudinal data, including years before baseline, are rare. METHODS: An observational single-center study was carried out. We analyzed data from 64 consecutive patients with multiple sclerosis. RESULTS: After 1 year of treatment ($n = 64$), score on the Expanded Disability Status Scale (EDSS) decreased by 0.47 points ($P = 0.047$) and the annualized relapse rate (ARR) decreased by 82% ($P < 0.001$). After 2 years ($n = 41$), EDSS score was still reduced by 0.28 (not significant) and ARR was reduced by 69% ($P < 0.001$). After 3 years ($n = 23$), EDSS score was reduced by 0.26 (not significant), and ARR was reduced by 77% ($P < 0.001$). Reduction of EDSS score and ARR did not depend on baseline ARR (1-2 vs >2) or EDSS score and was not biased by exceptional high disease activity or relapses around baseline. CONCLUSIONS: These real-life data reinforce that natalizumab is effective over years, reduces ARR, and stabilizes EDSS score independent of baseline ARR, baseline EDSS score, or baseline treatment.

DOI: <https://doi.org/10.1097/WNF.0b013e31824644e6>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-64660>

Journal Article

Accepted Version

Originally published at:

Kallweit, U; Jelcic, I; Braun, N; Fischer, H; Zörner, B; Schreiner, B; Sokolov, A A; Martin, R; Weller, M; Linnebank, M (2012). Sustained efficacy of natalizumab in the treatment of relapsing-remitting multiple sclerosis independent of disease activity and disability at baseline: real-life data from a Swiss cohort. *Clinical Neuropharmacology*, 35(2):77-80.

DOI: <https://doi.org/10.1097/WNF.0b013e31824644e6>

Sustained efficacy of Natalizumab in the treatment of relapsing-remitting multiple sclerosis independent of disease activity and disability at baseline: real-life data from a Swiss cohort

Ulf Kallweit, MD; Ilijas Jelcic, MD; Nathalie Braun, MD, PhD; Heike Fischer, MD; Björn Zörner, MD; Bettina Schreiner, MD; Arseny A. Sokolov, MD; Roland Martin, MD; Michael Weller, MD; Michael Linnebank, MD

¹Department of Neurology, University Hospital Zurich, Frauenklinikstrasse 26, 8091 Zurich, Switzerland

Correspondence:

PD Dr. Michael Linnebank, Department of Neurology, University Hospital Zurich, Tel. +41 44 255 1111. Fax. +41 44 255 4380. Email: michael.linnebank@usz.ch

Running title: Natalizumab: Swiss real-life data

Key-words: Natalizumab, multiple sclerosis, EDSS, annual relapse rate, real-life

Word count: 1583

Abstract

Background:

Therapy of relapsing-remitting multiple sclerosis with Natalizumab (Tysabri, Biogen Idec) has been shown to be effective in the reduction of the clinical relapse rate and disability progression. However, real-life, longitudinal data, including years before baseline, are rare.

Methods:

An observational, single-center study was carried out. We analyzed data from 64 consecutive MS patients.

Results:

After one year of treatment (n=64), EDSS decreased by 0.47 points (p=0.047), the annualized relapse rate (ARR) by 82% (p<0.001). After two years (n=41), EDSS was still reduced by 0.28 (n.s.) and ARR by 69% (p<0.001). After three years (n=23), EDSS was reduced by 0.26 (n.s.), ARR by 77% (p<0.001). EDSS and ARR reduction did not depend on baseline ARR (1-2 vs. >2) or EDSS and were not biased by exceptional high disease activity or relapses around baseline.

Conclusion:

These real life data reinforce that natalizumab is effective over years, reduces ARR and stabilizes EDSS independent of baseline ARR, baseline EDSS or baseline treatment.

Kommentar [UK1]: Ich finde, dass gehört nicht zwingend ins abstract

Introduction

In the pivotal trial of natalizumab therapy of relapsing-remitting multiple sclerosis (RRMS; AFFIRM) [1], the clinical relapse rate was reduced by 68% compared with placebo in the first year of treatment. Furthermore, the probability of disability progression was decreased [1]. Several subsequent studies have confirmed the efficacy of natalizumab [2-9]. However, real-life longitudinal data, including years before baseline and especially on patients with a high EDSS at baseline, are rare. The present study provides such real-life data on 64 natalizumab-treated MS patients with a mean baseline EDSS of 4.5 ± 1.7 .

Patients and Methods

We conducted our study in compliance with the principles of the Declaration of Helsinki. An observational, monocenter study was carried out. Inclusion criteria were natalizumab treatment of RRMS according to the modified McDonald criteria [10] and availability of standardized clinical data, collected in our MS center, on at least one year before and one year after initiation of natalizumab treatment. Relapses were defined according to McDonald criteria. PASW statistics version 18 was used for ANOVA statistics (IBM, NY). When data were only available for a subset of patients, e.g. the 2y and 3y follow-up data, we compared the available data with those patients' baseline data only and excluded baseline data of the other patients. Data were exploratively stratified by baseline EDSS, baseline ARR, median disease duration and prior therapies.

Results

Sixty-four consecutive patients were enrolled. Two additional patients developed persisting anti-natalizumab-antibodies, one developed an allergic reaction and one developed a progressive multifocal leukoencephalopathy (PML). These four patients were not included into further analyses. At baseline, mean age ± 1 standard deviation was 41 ± 10 yr. (range: 22-68). Forty-seven patients (0.73) were female, 17 (0.27) were male. Disease duration was 9.9 ± 7.5 yr. (range: 0-29). Sixteen (0.25) patients had no prior immunomodulating or

immunosuppressive therapy other than steroids, 33 (52%) had been treated with beta-interferons only, 2 (3%) with glatiramer acetate only, 6 (9%) with both beta-interferons and glatiramer acetate one after another, and 7 (11%) had immunosuppressant treatment after beta-interferon and/ or glatiramer acetate treatment. Two patients had had an additional treatment with immunoglobulins.

Initiation of natalizumab treatment reduced the ARR in the entire population as well as in subgroups defined by a moderate or high ARR at baseline, respectively (*Table 1, Figure 1a*). The ARR was significantly reduced over the three years observed. In addition, natalizumab treatment also reduced the EDSS in the first year (*Table 1*). EDSS after two and three years was still lower in comparison to the baseline EDSS of the respective patients, but this was not significant, consistent with the smaller effect size and the smaller sample group. EDSS reduction was similar when stratified by limitation versus non-limitation of walking distance (*Figure 1b*). ARR and EDSS reduction were also independent of age, gender, or disease duration (not shown). Concerning patients without previous treatment, ARR baseline versus first year was 2.52 ± 0.87 versus 0.38 ± 0.74 (=85% reduction; $p < 0.001$). For patients receiving therapy with interferons or glatiramer acetate before the time of natalizumab initiation, ARR baseline versus first year was 1.95 ± 1.0 versus 0.29 ± 0.46 (=85% reduction; $p < 0.001$). In the first year of natalizumab treatment, EDSS improved in 22/64 (0.34) of patients and was stable in further 31/64 (0.48) patients. After two years, EDSS was better than baseline in 12/37 (0.32) and equal in 18/37 (0.49), and, after three years, EDSS was better in 6/23 (0.26) and equal in 13/23 (0.57). Clinical freedom of disease activity, i.e. no relapses and improved or stable EDSS, was observed in 43/59 (0.73), 13/37 (0.35), and 8/23 (0.35) patients after one, two and three years of natalizumab treatment. Regarding the years before baseline, ARR in the first year of treatment was significantly lower in comparison to one ($p = 0.012$) and two ($p = 0.001$) years before baseline, and also ARR of the second and third year after initiation was lower compared with one ($p = 0.017$ and $p = 0.010$) and two years ($p = 0.042$ and

Kommentar [UK2]: Sollen wir hier noch dazu schreiben („data not shown“)? Wenn wir noch eine Tabelle machen oder es in die Tabelle einfügen wird sie noch unübersichtlicher.
LASS ERSTMAL SO

$p=0.022$) before treatment initiation (*Table 1, Figure 1a*). This was not significant for EDSS (*Table 1*).

Discussion

Natalizumab has been shown to be effective in the treatment of RRMS [1-9]. However, longitudinal real-life data are rare. The Swedish national post-marketing surveillance study demonstrated sustained efficacy of natalizumab treatment for the Swedish population [9]. For the present population, similar results were obtained with data collected in our MS-center standardized to the McDonald criteria [10].

Natalizumab treatment is often initiated after a period of active disease resulting in a high baseline ARR. It cannot be excluded that, in the natural course of disease, the ARR of the next year would have been lower again, even without initiation of natalizumab therapy. Natural regression to the mean disease activity might be an important bias in respective studies using baseline data for the analysis of natalizumab efficacy, as Natalizumab often is initiated in a periods of high disease activity. Thus, comparison of follow-up data during Natalizumab therapy should not only been performed with baseline, but also with the disease activity before baseline. In our study, the ARR in the first three years was significantly lower not only in comparison to baseline data, but also to one and two years before baseline, demonstrating a sustained efficacy not derived from a bias concerning the evaluation of baseline data. Moreover, ARR reduction in the first year was identical in patients without treatment and patients under treatment with interferons or glatiramer acetate at treatment initiation. These data confirm the efficacy of natalizumab in patients with active disease, regardless if treatment naïve/ currently untreated or under first-line immunomodulating therapy. In our group of patients, the efficacy of natalizumab over three years according to the reduction on ARR and, in part, EDSS, was confirmed. The effect of natalizumab on patients with more advanced disease severity, i.e. with a reduced walking distance, was comparable to patients without reduced walking distance showing a remarkable stabilization

Kommentar [ML3]: immer gross oder klein

and slight improvement on EDSS indicating the positive effect of Tysabri on EDSS compared to the natural course of the disease. [11]

Conclusion

In conclusion, our study reinforces that natalizumab is efficient in the treatment of RRMS independent of baseline ARR, baseline EDSS or baseline treatment. Improvement after baseline is not likely to be biased by exceptionally high disease activity at the time-point of Tysabri initiation as shown by the data from year(s) before baseline. Efficacy of natalizumab therapy persists over years. The benefits of natalizumab treatment must be weighed against the risks.

References

1. Havrdova E, Galetta S, Hutchinson M, et al. Effect of... AFFIRM
2. Polman CH, O'Connor PW, Havrdova E, et al. A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *New England Journal of Medicine* 2006;354:899–910.
3. Prosperini L, Borriello G, Fubelli F, Marinelli F and Pozzilli C. Natalizumab treatment in multiple sclerosis: the experience of S.Andrea MS Centre in Rome. *Journal of the Neurological Science* 2010, Jun10. [Epub ahead of print]. PMID: 20535513
4. Oturai AB, Koch-Henriksen N, Petersen T, Jensen PE, Sellebjerg F and Sorensen PS. Efficacy of natalizumab in multiple sclerosis patients with high disease activity: a Danish nationwide study. *European Journal of Neurology* 2009; 16:420–423
5. Putzki N, Yaldizli O, Buhler R, Schwegler G, Curtius D and Tettenborn B. Natalizumab reduces clinical and MRI activity in multiple sclerosis patients with high disease activity: results from a multicenter study in Switzerland. *European Neurology* 2010; 63:101–106.

6. Memon A, Caon C, Perumal J, et al. Natalizumab is effective in RRMS patients who do not respond to sequential therapy with interferon beta and glatiramer acetate: results of a switching study. *Neurology* 2010, 74:A542
7. Fernandez O, Alvarenga MP, Guerrero M, et al. The efficacy of natalizumab in patients with multiple sclerosis according to level of disability: results of an observational study. *Multiple Sclerosis Journal* 2011,17:192-197
8. Kaufman MD, Lee R, Norton HJ. Course of relapsing-remitting multiple sclerosis before, during and after natalizumab. *Multiple Sclerosis Journal* 2010, Dec 6, [Epub ahead of print], PMID: 21135017
9. Miller DH, Khan OA, Sheremata WA, et al. A controlled trial of natalizumab for relapsing-multiple sclerosis. *New England Journal of Medicine* 2003;348:15–23.
10. Holmén C, Piehl F, Hillert J, et al. Swedish national post-marketing surveillance study of natalizumab treatment in multiple sclerosis. *Multiple Sclerosis Journal* 2011, Jan 12. [Epub ahead of print] PubMed PMID: 21228027.
11. Polman C, Reingold S, Edan G., et al. Diagnostic criteria for multiple sclerosis: 2005 revisions to the “McDonald Criteria”. *Annals of Neurology* 2005;58:840-846
12. Confavreux C, Vukusic S, Moreau T, et al. Relapses and progression of disability in multiple sclerosis. *N Engl J Med*. 2000;343:1430-1438.

Kommentar [ML4]: im text nur bis ref 11, ist da was verschoben?

Figure title and legend:

Figure 1a: Annualized relapse rate

Annualized relapse rate (ARR) before and after initiation of natalizumab treatment. * $p < 0.05$, *** $p < 0.001$

Figure 1b: EDSS

EDSS differences from baseline ($t=0$: EDSS=4.52) beginning two years before to three years after therapy. * $p < 0.05$, ** $p < 0.01$

Funding

- There was no funding or sponsorship for this study.

Disclosure of conflicts of interest

- UK received honoraria from serving on an international expert panel meeting from Merck Serono.
- IJ, NB, HF, BZ, BS, AAS report no disclosures.
- MW received honoraria for advisory boards and research support from Merck Serono
- ML received grants from Bayer Health Care, Biogen Idec, Merck Serono, Novartis and Teva Pharmaceuticals. He has received personal compensation for participating in advisory boards and acting as consultant for Bayer Health Care, Biogen Idec, Merck-Serono, Novartis and Teva Pharmaceuticals.